

Rutgers University, New Jersey Agricultural Experiment Station

2013 Potato Disease Forecasting Report

July 12, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
7/8	1	45	512	0	57	491	0	19	507	0	86	504	0	24	480
7/9	0	45	521	0	57	498	0	19	514	1	87	512	1	25	487
7/10	2	47	529	0	57	505	0	19	519	2	89	517	0	25	493
7/11	0	47	537	0	57	511	0	19	526	1	90	524	0	25	502

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>