

Rutgers University, New Jersey Agricultural Experiment Station

2013 Potato Disease Forecasting Report

July 16, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
7/11	0	47	537	0	57	511	0	19	526	1	90	524	0	25	502
7/12	2	49	547	0	57	521	2	21	535	2	92	534	5	30	511
7/13	2	51	556	2	59	528	2	23	542	3	95	540	2	32	517
7/14	2	53	562	2	61	532	1	24	547	3	98	545	1	33	525
7/15	0	53	568	0	61	538	0	24	552	1	99	549	0	33	531

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>