

# Rutgers University, New Jersey Agricultural Experiment Station

## 2013 Potato Disease Forecasting Report

### July 23, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
7/18	0	53	582	0	61	550	0	24	563	0	100	560	0	33	545
7/19	0	53	586	0	61	553	0	24	566	2	102	562	0	33	548
7/20	0	53	592	0	61	556	0	24	571	0	102	566	0	33	550
7/21	0	53	600	0	61	562	0	24	576	2	104	572	0	33	556
7/22	0	53	608	2	63	569	0	24	583	4	108	578	1	34	563

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>