

# Rutgers University, New Jersey Agricultural Experiment Station

## 2013 Potato Disease Forecasting Report

### July 26, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
7/22	0	53	608	2	63	569	0	24	583	4	108	578	1	34	563
7/23	3	56	617	3	66	576	1	25	589	3	111	585	1	35	571
7/24	0	56	657	0	66	585	0	25	598	1	112	594	1	36	578
7/25	0	56	635	0	66	595	0	25	608	0	112	603	0	36	588

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>