

Rutgers University, New Jersey Agricultural Experiment Station

2013 Potato Disease Forecasting Report

July 30, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
7/25	0	56	635	0	66	595	0	25	608	0	112	603	0	36	588
7/26	0	56	644	0	66	603	0	25	617	2	114	612	0	36	597
7/27	0	56	653	1	67	611	0	25	625	2	116	621	1	37	506
7/28	0	56	663	1	68	620	0	25	634	3	119	629	1	38	515
7/29	2	58	673	2	70	629	2	27	643	3	122	638	1	39	524

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>