

Rutgers University, New Jersey Agricultural Experiment Station

2013 Potato Disease Forecasting Report

August 2, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
7/29	2	58	673	2	70	629	2	27	643	3	122	638	1	39	624
7/30	0	58	681	0	70	637	0	27	652	1	123	646	0	39	633
7/31	0	58	690	1	71	646	0	27	661	1	124	655	0	39	641
8/1	0	58	700	0	71	656	0	27	671	0	124	665	0	39	651

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>