

# Rutgers University, New Jersey Agricultural Experiment Station

## 2013 Potato Disease Forecasting Report

### August 22, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
8/18	0	71	853	1	80	801	0	34	817	1	150	808	0	51	804
8/19	2	73	863	2	82	810	0	34	827	2	152	817	0	51	813
8/20	0	73	872	0	82	820	0	34	836	2	154	825	0	51	822
8/21	0	73	879	0	82	826	0	34	842	1	155	832	0	51	830

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>