

# Rutgers University, New Jersey Agricultural Experiment Station

## 2013 Potato Disease Forecasting Report

### August 27, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
8/21	0	73	879	0	82	826	0	34	842	1	155	832	0	51	830
8/22	2	75	888	2	84	834	0	34	850	2	157	841	0	51	838
8/23	0	75	898	0	84	843	0	34	860	4	161	850	0	51	847
8/24	0	75	907	0	84	852	0	34	869	1	162	859	0	51	856
8/25	0	75	916	0	84	860	0	34	877	1	163	867	0	51	863
8/26	0	75	925	0	84	868	0	34	886	1	164	876	0	51	872

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>