

Rutgers University, New Jersey Agricultural Experiment Station

2013 Potato Disease Forecasting Report

August 30, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
8/26	0	75	925	0	84	868	0	34	886	1	164	876	0	51	872
8/27	0	75	933	0	84	875	0	34	893	0	164	882	0	51	879
8/28	0	75	943	0	84	885	0	34	902	5	169	892	0	51	888
8/29	2	77	952	1	85	894	1	35	911	4	173	901	0	51	896

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>