

# Rutgers University, New Jersey Agricultural Experiment Station

## 2013 Potato Disease Forecasting Report

### August 6, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
8/1	0	58	700	0	71	656	0	27	671	0	124	665	0	39	651
8/2	5	63	709	4	75	664	2	29	679	6	130	672	4	43	660
8/3	0	63	719	0	75	674	0	29	689	1	131	682	0	43	670
8/4	0	63	729	0	75	683	0	29	698	1	132	691	0	43	680
8/5	0	63	737	0	75	692	0	29	707	0	132	700	0	43	688

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>