

Rutgers University, New Jersey Agricultural Experiment Station

2013 Potato Disease Forecasting Report

August 9, 2013

A. Wyenandt, W. Kline, J. Ingerson-Mahar

Date	Pittstown			Hamilton			Upper Deerfield			Mount Holly			Woodbine		
	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days	Daily DSV	Total DSV	Total P-days
8/5	0	63	737	0	75	692	0	29	707	0	132	700	0	43	688
8/6	0	65	746	0	75	700	0	29	716	1	133	709	0	43	697
8/7	0	65	756	0	75	710	0	29	726	2	135	719	2	45	706
8/8	2	67	764	2	77	717	0	29	734	4	139	725	0	45	716

- Late blight was found on potato and processing tomato in Salem county on 7/2 and in organic tomato in Mercer County. Late blight has also been detected in DE, VA and MD in recent weeks. Scouting on a regular basis is highly recommended. All potato and tomato growers who have not initiated a standard protectant fungicide program should do so. All growers should also consider adding a late blight specific fungicide to their programs as long as current weather patterns continue.
- All late blight development updates are being posted under the [Vegetable Crop section of the Plant and Pest Advisory](#).
- To track late blight in the US please visit <http://usablight.org/>