



Section 18 Exemption

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision, and only for the uses covered by the certified applicator's certification.

For distribution and use only in New Jersey under an emergency exemption authorized under Section 18 of FIFRA

EPA FILE SYMBOLS: 13-NJ-01, 13-NJ-02, 13-NJ-03

Bifenture[®] EC Agricultural Insecticide

EPA Reg. No. 70506-57

All applicable directions, restrictions, and precautions on the EPA registered product labels as well as those on these directions for use must be followed. These directions for use must be in the possession of the user at the time of pesticide application. Any adverse effects resulting from the use of this product under this emergency exemption must be immediately reported to the New Jersey Department of Environmental Protections.

This exemption is effective July 16, 2013 through October 15, 2013.

APPLES, PEACHES, NECTARINES

PEST CONTROLLED	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	PHI (days)
Brown Marmorated Stink Bug (<i>Halyomorpha halys</i>)	5.12-12.8 fl. ozs./A (0.08-0.20 lbs. ai)	By Ground Only: Apply as a dilute spray (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre). Use higher rates under heavy insect pressure. For best control thorough coverage is necessary. Apply as necessary to maintain control using a minimum of 30-day spray intervals. Applications should be applied when populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area. Restrictions: Do not apply more than 32 fl. ozs. (0.50 lbs ai) per acre per season. For all bifenthrin products used, do not apply more than a total of 0.50 lbs. ai/acre per season. Do not apply this product until after petal fall. Do not allow entry into treated areas for 12 hours following application (REI = 12 hours). Do not graze livestock in treated orchards or cut treated cover crops for feed.	14