
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1388A-S1 1114

This supplemental label expires on 11/20/2017 and must not be used or distributed after this date.

Micora™

Fungicide

For control of diseases caused by Downy Mildew:

- Basil grown for transplant and retail sale to consumers

Active Ingredient:

Mandipropamid*	23.3%
<i>Other Ingredients:</i>	76.7%
<hr/>	
Total:	100.0%

*CAS No. 374726-62-2

KEEP OUT OF REACH OF CHILDREN.

EPA Reg. No. 100-1388

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Micora™ as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

USE DIRECTIONS FOR BASIL TRANSPLANTS GROWN IN ENCLOSED GREENHOUSES WITH PERMANENT FLOORING FOR RE-SALE TO CONSUMERS

Crop	Disease	Rate fl oz/Acre (fl oz/5,000 sq ft)	Remarks
Basil	Downy mildew (<i>Peronospora belbahrii</i>)	8.0 (0.9)	<p>Apply no more than two applications containing Micora per crop before switching to a non-group 40 fungicide.</p> <p>One application may be made during the plug growing stage and one application may be made during the crop finishing stage.</p> <p>Begin applications prior to disease development and continue on a 7-10 day interval. Use the shorter interval under high pressure or when conditions are conducive to disease. Micora may be tank mixed with another fungicide labeled for downy mildew that has a different mode of action.</p> <p>The addition of a spreading/penetrating type adjuvant such as a non-ionic based surfactant or crop oil concentrate or blend is recommended.</p> <p>For best results, use sufficient water volume to provide thorough coverage. Micora may be applied by ground application or chemigation.</p>
<p>Specific Use Restrictions:</p> <ul style="list-style-type: none"> • Do not apply within 1 day of harvest or shipping (1-day PHI). 			

©2015 Syngenta

Micora™ is a trademark of a Syngenta Group Company