GROUP 4A INSECTICIDE

SECTION 18 EMERGENCY EXEMPTION

FOR USE AND DISTRIBUTION ONLY WITHIN THE STATE OF NEW JERSEY

FOR CONTROL OF BROWN MARMORATED STINK BUGS IN POME AND STONE FRUIT.



EFFECTIVE DATE: 8-17-2016 EXPIRATION DATE: 10-15-2016

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, Worker Protection Standard (WPS) requirements, and precautions on the federally registered product label for Scorpion 35 SL Insecticide (EPA Reg. No. 10163-317), as well as the Section 18 directions, must be followed.
- This label must be in the possession of the user at the time of pesticide application.
- Do not use Scorpion 35 SL Insecticide on pome and stone fruit after the expiration date of 10-15-2016



Do not apply this product while bees are foraging. This compound is highly toxic to honey bees. The persistence of residues and potential residual toxicity of dinotefuran in nectar and pollen suggests the possibility of chronic toxic risk to honey bee larvae and the eventual instability of the hive. This product is toxic to bees exposed to residues for more than 38 hours following treatment.

DIRECTIONS FOR USE

Direction for use on Pome and Stone Fruits			
PEST	RATE FL. OZ./A	REMARKS	RESTRICTIONS
Brown Marmorated Stink Bug	8 - 12 fl oz/A (0.203 lbs - 0.304 lbs ai/A)	Use adequate spray volumes to insure complete coverage. Higher water volumes may provide improved insect control. Begin application when field sampling indicates pest activity. Use higher label rates when pest pressure increases and especially when adults are actively migrating into the orchard. Repeat as needed to maintain control, but not more often than every 7 days.	Restriction: Do not apply more than a total of 24 fl oz/A of SCORPION 35SL INSECTICIDE (0.608 lb ai/A) per season. Do not make more than 2 applications per acre per season of this product or other products containing dinotefuran. Do not apply this product until after petal fall. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
		SCORPION 35SL INSECTICIDE may be tank mixed and/or alternated with other approved insecticides. To reduce the potential for resistance, use a recommended IPM program consisting of varying modes of action.	 Foliar Application Only Apply with ground equipment in adequate water to obtain uniform coverage (For best results, use at least 50 gallons/acre for ground applications). Do not apply SCORPION 35SL INSECTICIDE within three (3) days of harvest.

Scorpion® is a registered trademark of Gowan Company, LLC.

EPA Reg. No. 10163-317

EPA File Symbol: 16-NJ-01 (pome fruit); 16-NJ-02 (stone fruit)

NJ Reg. No. SECTION18-2016-NJ-01 (pome fruit) NJ Reg. No. SECTION18-2016-NJ-02 (stone fruit)



Gowan Company P.O. BOX 5569 Yuma, AZ 85366-5569